Volume 4, Number 6

August 2011



# SAGE WINDS

NATIONAL WEATHER SERVICE - BOISE, ID

http://www.weather.gov/boise


**AUGUST 2011** 

## We're Having a Heat Wave...

A massive ridge of high pressure has dominated the longwave weather pattern for over two weeks, causing abnormally hot and humid conditions across much of the Midwest and Southern Plains. Due to the positioning of this quasi-stationary high, Texas, Oklahoma, Missouri, Kansas, and Arkansas have experienced the brunt of the sweltering conditions. The water vapor image at top right has been overlayed with 500 mb heights to show the position of this system on July 18th.


The image at bottom left gives a sampling of high temperatures from the same afternoon.

Many record highs have been broken



across the country, including record high maximum temperatures *AND* record high minimum temperatures. In fact, 177 all-time records (highs and lows) were broken in the seven days spanning July 19th through July 25th. Refer to

the "Recent Records Totals" (below, courtesy of NCDC) for more details on recently broken all-time records.



ed by NOAA National Centers for Enviromental Prediction, Hydrometeorological Prediction Center

#### **Recent Records Totals**

	Daily   Monthly   All-Tin					
Period		Hi Max	Hi Min			
Yesterday	Jul 25, 2011	1	4			
Last 7 Days	Jul 19 - 25, 2011	58	119			
Last 30 Days	Jun 28 - Jul 25, 2011	96	183			
Last 365 Days	Jul 26, 2010 - Jul 25, 2011	141	357			
Month to Date	Jul 1 - 25, 2011	70	180			
Year to Date	Jan 1 - Jul 25, 2011	121	220			

### What is the "Heat Index"?

The "heat index" integrates standard air temperature and relative humidity to calculate an associated "human-perceived" temperature. The table to the right displays different temperatures, relative humidities, and their corresponding heat indices or "feels like" temperatures. Note that the relative humidity required to increase the heat index above the air temperature is much *less* at high temperatures than it is at cool temperatures.

									AIR	TEM	PER/	TUR	E (F)							
		80	82	84	86	88	90	92	94	96	98	100	102	104	106	108	110	112	114	116
	5	78	80	81	83	85	86	88	89	91	93	94	96	97	99	101	102	104	105	107
	10	78	80	81	83	84	86	88	89	91	93	95	97	99	101	103	105	107	109	111
	15	78	80	81	83	84	86	88	90	92	94	96	98	100	103	105	108	111	113	116
L	20	79	80	81	83	85	86	88	90	93	95	97	100	103	106	109	112	115	119	122
	25	79	80	82	83	85	87	89	91	94	97	100	103	106	109	113	117	121	125	129
ı	§ 30	79	80	82	84	86	88	90	93	96	99	102	106	110	114	118	122	127	132	137
		80	81	83	85	87	89	92	95	98	102	106	110	114	119	123	129	134	140	146
I	RELATIVE HUMIDITY 52 09 59 50 54 05 54 05 59 59 59 59 59 59 59 59 59 59 59 59 59	80	81	83	85	88	91	94	97	101	105	109	114	119	124	130	136	142	148	
ı	<b>월 45</b>	80	82	84	87	89	92	96	100	104	109	114	119	124	130	137	143			
ı	5 50	81	83	85	88	91	95	99	103	108	113	118	124	131	137					
ı	T 55	81	84	86	89	93	97	101	106	112	117	124	130	137						
ı	≦ 60	82	84	88	91	95	100	105	110	116	123	129								
I	65	82	85	89	93	98	103	108	114	121	128	136								
ľ	급 70	83	86	90	95	100	106	112	119	126										
ı		84	88	92	97	103	109	116	124											
	80	84	89	94	100	106	113	121												
	85	85	90	96	102	110	117													
	90	86	91	98	105	113					Blank	( area	a is w	here	dewp	oint:	> 85F			
	95	86	93	100	108															
	100	87	95	103																
Heat disorder risk CA							CAUTION EXTREME CAUTION DANGER EXTREME DANGER													

<<<

July 20, 2011:	<b>Observations &amp; Associated Heat Indices</b>
----------------	---

Location	Temp.	Dewpt.	Rel.	Heat In-	
			Humid- ity	dex	
Tulsa	103°	82°	52%	130°	
Des Moines	99 °	79°	53%	118°	
Omaha	100°	82°	57%	120°	
St. Louis	100°	82°	57%	120°	
Boise	82°	54°	38%	81°	

The table to the left gives a sampling of surface observations from several cities on July 20, 2011. Heat indices of 105°-130° F create dangerous conditions in which heat cramps and exhaustion are likely. At heat indices of 130° F and above, heat *stroke* becomes likely with continued exposure. Note Tulsa, Oklahoma's extremely dangerous heat index of 130° F.

"Preparing for Extreme Heat" - The Center for Disease Control

#### Weather In the News

RARE Snow in Atacama Desert, Chile (NASA)

Record Rainfall in Chicago (Chicago Tribune)

Tropical Storm Emily Barrels Toward Fragile Haiti (AFP)

Red Cross Offers Relief Aid to N. Korea (AP)

#### National Weather Service Boise Forecast Office

3833 S. Development Ave, Bldg 3807 Boise, Idaho, 83705

(208) 334-9860

Email us at: boi.spotter@noaa.gov

# The Phoenix Dust Storm—July 2011




Strong downburst winds of 50+ mph combined with pre-existing drought conditions to kick up one of the most impressive U.S. dust storms seen in years. The outflow winds initiated from strong thunderstorms located southeast of Phoenix, near Tucson, Arizona. The drought conditions allowed these outflow winds to pick up excess dust, sand, and loose topsoil, and subsequently carry it northwest into Phoenix.

In terms of size, the dust storm was immense. At its peak, it was estimated to be 1 mile high and approximately 100 miles wide along the leading edge. On the ground, it traveled a distance of more than 150 miles.

A detailed account of this storm, with video and additional photos, has been archived on the NWS—Phoenix website: "Major Dust Storm Moves Through Arizona".

Mike Olbinski, Phoenix photographer, captured some amazing video and stills of the dust storm. They can be viewed here: The Phoenix Haboob of July 5th, 2011—Mike Olbinski.



^^ Arizona Drought Conditions as of June 28, 2011— one week before the dust storm.

